

Unclassified

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE

## DTIC DOCUMENTATION PAGE

AD-A198 571

ELECTE

10 1988

1b RESTRICTIVE MARKINGS

3 DISTRIBUTION/AVAILABILITY OF REPORT

Approved for public release; distribution unlimited.

4 PERFORMING ORGANIZATION REPORT NUMBER(S)  
Technical Report No. DU/DC/TR-09

5 MONITORING ORGANIZATION REPORT NUMBER(S)

6a NAME OF PERFORMING ORGANIZATION  
Department of Chemistry  
Duke University6b OFFICE SYMBOL  
(if applicable)7a NAME OF MONITORING ORGANIZATION  
Office of Naval Research

6c ADDRESS (City, State, and ZIP Code)

Durham, NC 27706

7b ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street  
Arlington, VA 22217-50008a NAME OF FUNDING/SPONSORING  
ORGANIZATION  
Office of Naval Research8b OFFICE SYMBOL  
(if applicable)

9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-83-K-0572 R&amp;T Code 4135008---02

8c ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street  
Arlington, VA 22217-5000

10 SOURCE OF FUNDING NUMBERS

PROGRAM  
ELEMENT NO  
NRPROJECT  
NO  
053TASK  
NO  
841WORK UNIT  
ACCESSION NO11 TITLE (Include Security Classification) The First Organogallium Four-Membered Ring Compound with Arsenic,  
Halogen Mixed Bridging: Synthesis and Crystal Structure of  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$ 12 PERSONAL AUTHOR(S)  
R.L. Wells, W.K. Holley, S. Shafieezad, A.T. McPhail, and C.G. Pitt13a TYPE OF REPORT  
Technical13b TIME COVERED  
FROM TO14 DATE OF REPORT (Year, Month, Day)  
88-08-0315 PAGE COUNT  
516 SUPPLEMENTARY NOTATION Accepted for presentation at the Fifth International Symposium on  
Inorganic Ring Systems, and for publication in Phosphorus and Sulfur.

17 COSATI CODES

FIELD

GROUP

SUB-GROUP

18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Organogallium-arsenic; Four-Membered Ring; Arsenic,  
Halogen Mixed Bridging; Synthesis; Crystal Structure.

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

See attached.

20 DISTRIBUTION/AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS21 ABSTRACT SECURITY CLASSIFICATION  
Unclassified22a NAME OF RESPONSIBLE INDIVIDUAL  
Richard L. Wells22b TELEPHONE (Include Area Code)  
(919)684-6404

22c OFFICE SYMBOL

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted

All other editions are obsolete

88 8 08 135

SECURITY CLASSIFICATION OF THIS PAGE

Unclassified

Attachment to DD Form 1473

Block Number 19. ABSTRACT

THE FIRST ORGANOGALLIUM FOUR-MEMBERED RING COMPOUND WITH  
ARSENIC, HALOGEN MIXED BRIDGING: SYNTHESIS AND CRYSTAL  
STRUCTURE OF  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$

RICHARD L. WELLS\*, WILLIAM K. HOLLEY, SOHEILA SHAFIEEZAD,  
ANDREW T. MCPHAIL, AND COLIN G. PITT  
Department of Chemistry, Paul M. Gross Chemical Laboratory,  
Duke University, Durham, NC 27706, U.S.A.

Abstract  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$  (1) has been isolated from the products of the room temperature reaction of  $\text{Ph}_2\text{GaCl}$  with  $(\text{Me}_3\text{Si})_3\text{As}$  (reactants mixed in both a 2:1 and a 3:1 mole ratio). A mixture of 1 and  $[\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2]_2$  (2) was isolated after heating a 1:1 mole ratio combination of the same reactants. Reaction of pure 2 [prepared from  $\text{Ph}_2\text{GaCl}$  and  $\text{LiAs}(\text{SiMe}_3)_2$ ] with  $\text{Ph}_2\text{GaCl}$  resulted in the formation of (1). Prolonged heating of 1 produced a mixture of 2,  $\text{Me}_3\text{SiCl}$  and unidentified products. Compound 1 was structurally characterized by a single-crystal X-ray analysis and shown to be the first organogallium four-membered ring compound with both an arsenic and a halogen bridge. The ring of 1 is clearly non-planar as evidenced by the fact that the Cl atom is displaced from the Ga-As-Ga' plane to yield a dihedral angle of  $8.8^\circ$  between the Ga-As-Ga' and Ga-Cl-Ga' planes. Various other features of the structure of 1 are discussed.

*Gallium Arsenide, 1971*

QUALITY  
INSPECTED  
2

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A-1	

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0572

R&T Code 4135008---02

Technical Report No. DU/DC/TR-09

The First Organogallium Four-Membered Ring Compound with Arsenic, Halogen  
Mixed Bridging: Synthesis and Crystal Structure of  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$

by

R. L. Wells, W. K. Holley, S. Shafieezad  
A. T. McPhail, and C. G. Pitt

Prepared for Presentation at the Fifth International  
Symposium on Inorganic Ring Systems and for  
Publication in Phosphorus and Sulfur

Duke University  
Department of Chemistry  
Durham, NC 27706

August 3, 1988

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

THE FIRST ORGANOGALLIUM FOUR-MEMBERED RING COMPOUND WITH  
ARSENIC, HALOGEN MIXED BRIDGING: SYNTHESIS AND CRYSTAL  
STRUCTURE OF  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$

RICHARD L. WELLS\*, WILLIAM K. HOLLEY, SOHEILA SHAFIEEZAD,  
ANDREW T. McPHAIL, AND COLIN G. PITT  
Department of Chemistry, Paul M. Gross Chemical Laboratory,  
Duke University, Durham, NC 27706, U.S.A.

Abstract  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$  (1) has been isolated from the products of the room temperature reaction of  $\text{Ph}_2\text{GaCl}$  with  $(\text{Me}_3\text{Si})_3\text{As}$  (reactants mixed in both a 2:1 and a 3:1 mole ratio). A mixture of 1 and  $[\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2]_2$  (2) was isolated after heating a 1:1 mole ratio combination of the same reactants. Reaction of pure 2 [prepared from  $\text{Ph}_2\text{GaCl}$  and  $\text{LiAs}(\text{SiMe}_3)_2$ ] with  $\text{Ph}_2\text{GaCl}$  resulted in the formation of (1). Prolonged heating of 1 produced a mixture of 2,  $\text{Me}_3\text{SiCl}$  and unidentified products. Compound 1 was structurally characterized by a single-crystal X-ray analysis and shown to be the first organogallium four-membered ring compound with both an arsenic and a halogen bridge. The ring of 1 is clearly non-planar as evidenced by the fact that the Cl atom is displaced from the Ga-As-Ga' plane to yield a dihedral angle of  $8.8^\circ$  between the Ga-As-Ga' and Ga-Cl-Ga' planes. Various other features of the structure of 1 are discussed.


INTRODUCTION

In gallium chemistry, four-membered ring formation is known to occur via bridging of gallium centers by two arsenic atoms<sup>1</sup> or two halogen atoms<sup>2</sup>, but the literature contains no references to this occurring through one of each of these atoms. Here we report the synthesis and crystal structure of  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$  (1), the first organogallium four-membered ring compound resulting from arsenic, halogen mixed bridging. We also report the synthesis of  $[\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2]_2$  (2). The fact that 1 can be prepared from  $\text{Ph}_2\text{GaCl}$  and  $(\text{Me}_3\text{Si})_3\text{As}$  again exemplifies the utility of

dehalosilylation between a silylarsine and a halogallane in preparing novel gallium-arsenic systems.<sup>1,3</sup>


#### RESULTS AND DISCUSSION

Combining  $C_6H_6$  solutions of  $Ph_2GaCl$ <sup>4</sup> and  $(Me_3Si)_3As$ <sup>5</sup> (2:1 mole ratio), followed by stirring at room temperature and removal of solvent and  $Me_3SiCl$ , gave a white solid. A ligroin extract of the solid afforded **1** as white crystals [mp 145-146 °C (dec), 55.3% yield]. A satisfactory molecular weight was obtained by cryoscopic measurements. NMR:  $^{13}C\{^1H\}$  ( $C_6D_6$ )  $\delta$  3.14 (s,  $Me_3Si$ ), 128.30, 128.72, 135.81, 146.57 (m, Ph).



Likewise, mixing solutions of  $Ph_2GaCl$  and  $(Me_3Si)_3As$  (3:1 mole ratio) resulted in the formation of **1** as the predominant product (53.5% yield).

A mixture of **1** and **2**,  $Me_3SiCl$  and other unidentified products were isolated after heating (76 °C) a 1:1 mole ratio combination of  $Ph_2GaCl$  and  $(Me_3Si)_3As$  in  $C_6H_6$ . Compounds **1** and **2** were identified as the components of the mixture by comparison of the  $^{13}C\{^1H\}$  NMR spectrum of the mixture with those of authentic samples of **1** and **2**. The latter was prepared from  $Ph_2GaCl$  and  $LiAs(SiMe_3)_2$ <sup>5</sup> (1:1 mole ratio) in benzene and isolated as white crystals [mp 229-230 °C (dec), 33.3% yield], and it was determined to be a dimer in solution by cryoscopic molecular weight measurements. NMR:  $^{13}C\{^1H\}$  ( $C_6D_6$ )  $\delta$  4.82 (s,  $Me_3Si$ ), 127.75, 128.02, 137.62, 149.55 (m, Ph).



The production of **1** was also accomplished by allowing  $Ph_2GaCl$  and **2** (2:1 mole ratio in  $C_6D_6$ ) to react in a sealed NMR tube.

# RING COMPOUND WITH ARSENIC, HALOGEN MIXED BRIDGING

Heating a sample of 1 in C<sub>6</sub>D<sub>6</sub> in a sealed NMR tube for one month at 80 °C resulted in the formation of 2, Me<sub>3</sub>SiCl and other unidentified products.

An X-ray crystal structure analysis of 1 revealed that the asymmetric unit comprises a discrete molecule (Figure 1) containing the heretofore unknown As- and Cl-bridged four-membered Ga-As-Ga'-Cl ring. That this ring is not strictly planar, and thus




FIGURE 1 Molecular structure of  $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$  (1) (hydrogen atoms omitted for clarity). Selected distances (Å) and angles (°): Ga-As 2.469(2), Ga'-As 2.463(2), Ga-Cl 2.412(3), Ga'-Cl 2.409(4), Si-As 2.359(4), Si'-As 2.367(4), Ga-As-Ga' 88.70(7), Ga-Cl-Ga' 91.3(1), As-Ga-Cl 89.5(1), As-Ga'-Cl 89.8(1), Si-As-Si' 111.0(2), C(11)-Ga-C(17) 120.0(5), C(11')-Ga'-C(17') 122.6(5).

the molecule deviates from exact  $C_{2v}$  symmetry presumably to relieve unfavorable non-bonded intramolecular interactions between substituents in such a symmetric form, is manifested by the Cl atom displacement of 0.256 Å from the Ga-As-Ga' plane (the associated angle between the Ga-As-Ga' and Ga-Cl-Ga' planes is  $8.8^\circ$  and the mean endocyclic dihedral angle about the ring bonds is  $6.2^\circ$ ). The extent of the departure from exact planarity in **1** is somewhat less than in the (Ga-As)<sub>2</sub> ring of  $\{[(Me_3SiCH_2)_2As]_3Ga\}_2$ <sup>6</sup> where more severe overcrowding of the bulkier ring substituents leads to corresponding interplanar and dihedral angles of  $13.6^\circ$  and  $10.2^\circ$ , respectively. In contrast to the situation in (Ga-As)<sub>2</sub> rings where the endocyclic angles subtended at As and Ga differ significantly [range:  $94.57(4) - 96.02(4)^\circ$  and  $83.58(4) - 85.02(2)^\circ$ , respectively],<sup>1</sup> those in **1** are almost equal [ $88.70(7)^\circ$  at As;  $89.5(1)$  and  $89.8(1)^\circ$  at Ga]. The bond angle at the bridging Cl atom [ $91.3(1)^\circ$ ] is nearly the same as that of  $91.4(1)^\circ$  in  $[Ga(C_5H_5)Cl_2]_2$ (**3**)<sup>7</sup> and lies in the middle of the range of  $86(2)^\circ$  in  $(GaCl_3)_2$ (**4**)<sup>8</sup> and the mean of  $97.4(2)^\circ$  in  $[Ga(C_5Me_5)_2Cl]_2$  (**5**).<sup>7</sup> The mean C-Ga-C angle at  $121.3(5)^\circ$  is close to the corresponding value of  $120.8(2)^\circ$  in  $[(Me_3SiCH_2)_2AsGaPh_2]_2$ .<sup>9</sup> A significantly larger Si-As-Si' angle [ $111.0(2)^\circ$ ] is present in **1** than in  $[(Me_3Si)_2AsLi \cdot DME]_2$  (DME = 1, 2-dimethoxyethane) (**6**) [ $103.2(4)^\circ$ ].<sup>10</sup> The mean Ga-As bond length in **1** [2.466(2) Å] is shorter than any found within (Ga-As)<sub>2</sub> rings [range: 2.513(1) - 2.581(1) Å],<sup>1</sup> whereas the mean Ga-Cl distance at 2.411(4) Å is longer than the corresponding length in **4** [2.29(9) Å] as well as the mean of those in **3** [2.363(3) Å], but it is shorter than the mean in **5** [2.448(7) Å]. The mean Si-As distance at 2.363(4) Å is significantly longer than in **6** [2.307(7) Å].

Finally, the <sup>13</sup>C NMR spectrum and the experimentally determined molecular weight of **1** indicate it has the same molecular structure in solution as in the solid state.

# RING COMPOUND WITH ARSENIC, HALOGEN MIXED BRIDGING

## ACKNOWLEDGEMENT

We thank the Office of Naval Research for financial support.

## REFERENCES

1. A. P. Purdy, R. L. Wells, A. T. McPhail, and C. G. Pitt, Organometallics, **6**, 2099 (1987); and references cited therein.
2. For a general discussion and examples, see M. J. Taylor, Comprehensive Coordination Chemistry, G. Wilkinson, Editor-in-Chief (Pergamon Press, Oxford, 1987), Vol. 3, Chap. 25.1.
3. R. L. Wells, S. Shafieezad, A. T. McPhail, and C. G. Pitt, J. Chem. Soc., Chem. Commun., 1823 (1987).
4. P. G. Perkins and M. E. Twentyman, J. Chem. Soc., 1038 (1965).
5. V. G. Becker, G. Gutekunst, and H. J. Wessely, Z. Anorg. Allg. Chem., **462**, 113 (1980).
6. R. L. Wells, A. P. Purdy, K. T. Higa, A. T. McPhail, and C. G. Pitt, J. Organomet. Chem., **325**, C7 (1987).
7. O. T. Beachley Jr., R. B. Hallock, H. M. Zhang, and J. L. Atwood, Organometallics, **4**, 1675 (1985).
8. S. C. Wallwork and I. J. Worrall, J. Chem. Soc., 1816 (1965).
9. R. L. Wells, A. P. Purdy, A. T. McPhail, and C. G. Pitt, J. Organomet. Chem., **308**, 281 (1986).
10. V. G. Becker and C. Witthauer, Z. Anorg. Allg. Chem., **492**, 28 (1982).



D/1113/87/2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 92132	1